

A Short Communication:

Cancer Mortality in a Chinese Population
Exposed to Hexavalent Chromium in Water

Prepared by:

Zhang, JinDong¹ and ShuKun Li²

¹ JinZhou Health and Anti-Epidemic Station (currently retired), JinZhou City, LiaoNing
Province, China; ~~Recently deceased~~

~~JinDong Zhang~~

~~2200103~~

Submitted to either:

Archives of Environmental Health

Environmental Health Perspectives

American Journal of Public Health

~~Environmental and Occupational Medicine~~

CONFIDENTIAL DRAFT

November 16, 1995

Abstract

This report is a clarification and further analysis of our previously published mortality study regarding an incident of groundwater contamination with hexavalent chromium in the JinZhou area of China ~~Wang et al. 1998~~.⁽¹⁾ At the beginning of the contamination episode in 1965, average concentrations of hexavalent chromium (Cr^{+6}) in the affected groundwater of villages ranged from 0.004 ppm to 2.6 ppm. To assess the potential effect on local cancer rates of ~~drinking~~ well water ~~containing~~ Cr^{+6} exposure for up to 7 years, we conducted a retrospective mortality study of approximately 100,000 residents living in selected regions and villages of the JinZhou suburbs during 1970-1978.⁽²⁾ In the ~~our previous report we stated that five~~ Cr^{+6} contaminated villages combined, a significant excess of overall cancer mortality was observed ($p = 0.04$), but individual village mortality rates were inversely correlated with the amount of Cr^{+6} contamination in well water. Further analysis revealed no clear statistical increase in cancer mortality in the three villages adjacent to the source ($p = 0.25$) where 57% of the wells exceeded the WHO safe drinking water limit of 0.05 ppm Cr^{+6} . ~~Further~~ In contrast, a ~~greater~~ more substantial excess of cancer mortality ($p = 0.10$) was found in the two most distant villages that were ~~most distant from the source of contamination~~ last to receive Cr^{+6} contaminated groundwater and that had groundwater Cr^{+6} concentrations below 0.05 ppm. These results do not indicate an association of cancer mortality with exposure to Cr^{+6} contaminated groundwater. The observed pattern of cancer mortality ~~observed in the two villages~~ might reflect the influence of lifestyle or environmental factors not related to Cr^{+6} ~~or other factors may be involved~~. Further follow-up of this cohort is recommended to assess the possible influence of Cr^{+6} and other risk factors on cancer mortality.

Background

The JinZhou area of LiaoNing Province is composed of a downtown area and six suburb regions: Nuer River Region, ZhongTun Region, GuoShu Region, West Suburb Region, North Suburb

Region, and XueJia Region (Figure 1). The suburb regions are primarily agricultural but are the home of several industrial plants. JinZhou Alloy Plant and No.6 Petroleum Plant are the two largest plants. JinZhou Alloy Plant started regular chromium production in 1965, at which time a large amount of Cr^{+6} containing waste water was discharged into the dry river bed.⁽¹⁾ The discharged waste water contributed to the Cr^{+6} contamination at the beginning of this episode via a shallow aquifer beneath a dry river bed. Later, leachate from chromite ore processing residues from the plant was the main source of the Cr^{+6} contamination by the same groundwater aquifer below the river bed. The spent Ore residue had been stockpiled uncovered next to the plant where precipitation caused the rapid dissolution and leaching of Cr^{+6} to the shallow groundwater aquifer used for drinking water. A long and narrow contaminated area was formed along the dry river bed of the Old Nuer River soon after the Alloy Plant began operations in 1965 (Figure 1). The contamination source was not fully controlled until 1982 when a seepage prevention wall was built around the ore residue dump site. Interim remediation measures included the addition of ferrous sulfate to the ore residues, which reduced the Cr^{+6} but also polluted the aquifer with sulfates at concentrations up to several hundred ppm. Residents living in the villages located along the Old Nuer River were exposed to Cr^{+6} by drinking using well water that had been contaminated with Cr^{+6} . The distribution of Cr^{+6} in the groundwater plume under the well water concentrations of Cr^{+6} investigated thoroughly in 1965 is shown for each of the five villages in Figure 2.

Retrospective Mortality Studies

We conducted a series of retrospective mortality studies of the approximately 100,000 residents living in the JinZhou suburb regions in 1970-1978. Most of the approximately 100,000 people in the five villages along the Old Nuer River in JinZhou Suburb Regions.⁽²⁾ Most residents were farmers (>95%) who had lived in the JinZhou suburb regions (>95%) for most of their lives. Residences were concentrated in agricultural villages, and there was minimal migration within the population. We examined the death records in the local police departments to locate all

deaths that occurred in this population between 1970 to 1978. A standard form was used to abstract the data and to identify record the cause of each death. All survey staff received training, and a follow-up survey was conducted for part of the death records to ensure the quality of the abstracted data. Age-adjusted cancer rates were calculated for each of the six regions and for each of the five villages using the contaminated groundwater in the contamination pathway. The death rate was calculated by dividing the observed number of cancer deaths in 1970-1978 by the age-adjusted population size total number of follow-up years. The total number of follow-up person-years is estimated as the product of the length of follow-up (9 years) and the estimated size of the population in 1975, the midpoint of the follow-up period. Village-specific all cancer mortality rates were statistically compared to rates for Liaoning Province using the Poisson distribution. (5, pg 69).

Groundwater Cr⁶⁺ contamination In 1965 Cr⁶⁺ was measured at drinking water wells for each village which draw from the contaminated aquifer in the contamination pathway (Table 1, Figure 2). In general, higher concentrations levels of the Cr⁶⁺ contamination occurred in the villages closer to the pollution source. Dose-response relationships were examined using the distance of each village from the source as a surrogate for exposure. The Poisson regression model was used ^{by transformation} wherein the expected rate of cancer was assumed to change in a linear manner based on depends linearly on distance from the contamination source: $\ln(\text{rate of cancer}) = a + (b \times \text{distance})$. Negative values for the slope b indicate that proximity to the source was associated with greater cancer death rates. (5, chapter 4).

Results

The adjusted cancer death rate for the six suburb regions are as follows: Nuer River Region, 68.8 per 100,000 people per year; for ZhongTun Region, 68.4; for GuoShu Region, 64.7; for West Suburb Region, 54.3; for XueJia Region, 57.5; for North Suburb Region, 45.9. The rates for the first three of the regions mentioned these regions are comparable to the 1973-1975 rate of

66.1 per 100,000 for LiaoNing province.

The death rates for total cancer, lung cancer and stomach cancer for each village in the Cr⁶ contamination path are presented in Table 1. When the total cancer mortality for all five villages combined is compared to that of LiaoNing province, a statistically significant excess was observed (2-sided p-value=0.04). However, none of the individual villages showed a statistically significant excess at a 2-sided p-value of 0.1. Further investigation of the statistical trends was performed by combining the three villages that were closest to the contamination source and that had frequent well water measurements (Figure 2) in excess of the World Health Organization safe drinking water standard of 0.05 ppm Cr⁶ (WHO, 1989). This guideline is more stringent than that used in (0.1 ppm).

Those three villages showed no significant excess of total cancer mortality compared to the province rates (p-value = 0.25), while the two more distant villages with well water concentrations all below 0.05 ppm had a more substantial increase in total cancer mortality (p-value = 0.10). Thus, it is apparent that the increased mortality rate was not due to the contaminated water.

No statistical comparisons to province mortality rates could be made for site-specific cancer rates due to the lack of appropriate rate information. Stomach cancer comprises a large proportion of the total cancer rate for these villages and for all of China (Table 1). As shown, in general, villages closer to the contamination source do not have higher cancer rates, while one of the least contaminated villages, Shilitai, had a substantially higher rate of mortality from stomach cancer. Lung cancer mortality rates were highest in the more distant villages with Cr⁶ contamination below the World Health Organization limit of 0.05 ppm. The Pearson and Kendall dose-response models indicated a nonsignificant ($p > 0.05$), weak positive correlation between cancer rates and the distance from the source, contrary to the expected direction of association if Cr⁶ contamination were the cause of the associated with higher cancer rates.

Discussion

In our preliminary report, we identified the higher general mortality rates for the six villages with exposure to Cr^{+6} in groundwater. The Cr^{+6} contamination followed long and narrow pathway that started near the JinZhou Alloy Plant in the Nuer River Region and extended to the West Suburb Region. In general, total cancer mortality was elevated significantly for the population living along the plume. Exposure to Cr^{+6} contamination was highest for the populations closest to the plant and lowest for the populations farthest from the plant. Our further analysis revealed that the cancer death rates for the six villages in the contaminated area were not correlated with the degree of exposure to Cr^{+6} and that the predominant excess occurred in the villages with relatively low Cr^{+6} magnitude of Cr^{+6} contamination. Neither stomach cancer nor lung cancer indicated a positive association with Cr^{+6} concentration in well water contamination. The absence of a dose-response relationship between cancer and Cr^{+6} clarifies our translation and interpretation of our previous observation publication.⁽¹⁾ Although Cr^{+6} contamination cannot be ruled out completely as the reason for the high cancer death rates in these villages, these results do not support such a relationship. The relatively short latency period (i.e., 13 years, 1965 to 1978) covered in this study limits the causative interpretation of these findings regarding cancer and Cr^{+6} contamination, although the number of person-years represented is substantial (approximately 99,000). A mortality study with a longer follow-up period would be worthwhile and is recommended. Nonetheless, these results suggest that lifestyle or environmental factors not related to the Cr^{+6} contamination are likely to be a source of the variation in cancer rates. Additional studies to identify these factors are also recommended.

References

- (1) Zhang JianDong, Li XiLin. Chromium Pollution of Soil and Water in JinZhou. Journal of Chinese Preventive Medicine 1987; 21:262-264.

- (2) Zhang JianDong, Li XiLin. Study of the Effect of Environmental Pollution in JinZhou Area on Residents' Health I. Mortality Analysis. JinZhou's Health and Anti-Epidemic Station, ~~JinZhou's Health and Anti-Epidemic Station, China~~ 1979.
- (3) Agency for Toxic Substances and Disease Registry. Toxicological Profile for Chromium. US Department of Health & Human Services, Public Health Service, TP-92/08, 1993, pg 57.
- (4) WHO. 1989. IPCS environmental health criteria - chromium, vol. 61. World Health Organization, International Programme on Chemical Safety.
- (5) Breslow, N.E., Day, N.E. Statistical Methods in Cancer Research. Volume II - The Design and Analysis of Cohort Studies. International Agency for Research on Cancer Lyon, 1987.

Insert the following section in the cover page of the letter to the editor:

Please Address Correspondence to:

Dr. JinDong Zhang

c/o Tony Ye

972 Riley Drive

Albany, California 94706

(510) 748-5675

Table 1


Well Water Hexavalent Chromium (Cr^{6+}) Concentrations and Cancer Mortality Death-Rate Information
for Villages ~~possible Contamination Pathway~~ in the Contamination Pathway

	Village					Range of Average Rates in China*	Dose-Response: Coefficient(p value)
	JinChangBao	Nuer River	Yang-Xing	Shi Li Tai	WenJiaTun		
Distance (km) from the JinZhou alloy plant	1.4	1.5	3.0	3.5	5.0	-	-
Average Cr^{6+} concentration in water wells in 1965	0.031 ppm (123 wells)	2.6 ppm (170 wells)	0.18 ppm (50 wells)	0.02 ppm (21 wells)	0.004 ppm (33 wells)	-	-
Frequency of wells exceeding 0.05 ppm	12%	95%	40%	0%	0%	-	-
<u>Cancer Death Rates per 100,000 for 1970-1978</u>							
All Cancer (p value)	83.6 (0.32)	71.9 (0.74)	76.8 (0.62)	93.0 (0.12)	91.1 (0.26)	29.8-102	0.05 (p=0.57)
Stomach Cancer	36.7	28.0	36.5	55.2	27.7	5.2-40.2	0.04 (p=0.74)
Lung Cancer	13.2	15.0	21.4	-	20.8	1.8-17.8	0.12 (p=0.59)
Person-years	26,179	24,792	9,703	23,225	14,950	-	-

* Range of the average cancer rates for the 30 provinces in China (National Cancer Control Office of China, 1979).

Figure 2

Distribution of Cr concentrations in well water of villages
in the JinZhou Area in 1965






FIGURE 1
 MAP OF JINZHOU AREA; INCLUDING THE
 CHROMIUM CONTAMINATED AREA

TY-0112